

Author Index

- Aggarwal, A., Merritt, Jr., J.W. and Nash, T.E.
Cysteine-rich variant surface proteins of *Giardia lamblia*, 39
- Aline, Jr., R.F., Scholler, J.K. and Stuart, K.
Transcripts from the co-transposed segment of variant surface glycoprotein genes are in *Trypanosoma brucei* polyribosomes, 169
- Anand, R., see Clark, J.T., 15
- Anders, R.F., see Favaloro, J.M., 297
- Åslund, L., see Franzén, L., 201
- Baginsky, W.F., see Schmatz, D.M., 263
- Barbet, A.F., Myler, P.J., Williams, R.O. and McGuire, T.C.
Shared surface epitopes among trypanosomes of the same serodeme expressing different variable surface glycoprotein genes, 191
- Beach, D.H., see Goad, L.J., 179
- Becker, I., see Ostoa-Saloma, P., 133
- Berens, R.L., see Goad, L.J., 179
- Bisby, R.M., see Browning, P.M., 57
- Blanton, R., see Reis, M.G., 113
- Brown, K.N., see Viriyakosol, S., 93
- Browning, P.M. and Bisby, R.M.
Qinghaosu does not affect the major thermotropic phase transition in model membranes of dipalmitoylphosphatidylcholine, 57
- Bülow, R., Nonnengässer, C. and Overath, P.
Release of the variant surface glycoprotein during differentiation of bloodstream to procyclic forms of *Trypanosoma brucei*, 85
- Button, L.L., Russell, D.G., Klein, H.L., Medina-Acosta, E., Karess, R.E. and McMaster, W.R.
Genes encoding the major surface glycoprotein in *Leishmania* are tandemly linked at a single chromosomal locus and are constitutively transcribed, 271
- Cabrera, N., see Ostoa-Saloma, P., 133
- Capron, A., see Dissous, C., 49
- Chambers, A.E., see Rodrigues, V., 7
- Chaudhri, M., see Rodrigues, V., 7
- Clark, J.T., Donachie, S., Anand, R., Wilson, C.F., Heidrich, H.-G. and McBride, J.S.
46-53 Kilodalton glycoprotein from the surface of *Plasmodium falciparum* merozoites, 15
- Coggins, J.R., see Jarroll, E.L., 121
- Coppel, R.L., see Favaloro, J.M., 297
- Coquelet, H., see Paindavoine, P., 61
- Crewther, P.E., see Favaloro, J.M., 297
- Danforth, H.D., see Jenkins, M.C., 153
- Davis, A.H., see Reis, M.G., 113
- Delves, C.J., Goman, M., Ridley, R.G., Matile, H., Lensen, T.H.W., Ponnudurai, T. and Scaife, J.G.
Identification of *Plasmodium falciparum*-infected mosquitoes using a probe containing repetitive DNA, 105
- Diffley, P. and Mama, K.
Fixed and temporary fluctuations in the cell cycle of monomorphic lines of *Trypanosoma brucei gambiense*, 1
- Dissous, C. and Capron, A.
Schistosoma mansoni and its intermediate host *Biomphalaria glabrata* express a common 39 kilodalton acidic protein, 49
- Donachie, S., see Clark, J.T., 15
- Erlandsen, S.L., see Jarroll, E.L., 121
- Favaloro, J.M., Marshall, V.M., Crewther, P.E., Coppel, R.L., Kemp, D.J. and Anders, R.F.
cDNA sequence predicting an octapeptide-repeat antigen of *Plasmodium falciparum* (Short Communication), 297
- Fetterer, R.H., see Jenkins, M.C., 153
- Franzén, L., Wählin, B., Wahlgren, M., Åslund, L., Perlmann, P., Wigzell, H. and Pettersson, U.
Enhancement or inhibition of *Plasmodium falciparum* erythrocyte reinvasion in vitro by antibodies to an asparagine rich protein, 201
- Garfinkel, L., see Huber, M., 285
- Gitler, C., see Huber, M., 285
- Goad, L.J., Berens, R.L., Marr, J.J., Beach, D.H. and Holz, Jr., G.G.
The activity of ketoconazole and other azoles against *Trypanosoma cruzi*: biochemistry and chemotherapeutic action in vitro, 179
- Goman, M., see Delves, C.J., 105
- Gregory, W.F., see Maizels, R.M., 213
- Heidrich, H.-G., see Clark, J.T., 15
- Holloway, S.P., see Hyde, J.E., 247
- Holz, Jr., G.G., see Goad, L.J., 179
- Hori, H., see Watanabe, J.-i., 163
- Huber, M., Koller, B., Gitler, C., Mirelman, D., Revel, M., Rozenblatt, S. and Garfinkel, L.
Entamoeba histolytica ribosomal RNA genes are carried on palindromic circular DNA molecules, 285
- Hundt, E., see Knapp, B., 73
- Hyde, J.E., Kelly, S.L., Holloway, S.P., Snewin, V.A. and Sims, P.F.G.
A general approach to isolating *Plasmodium falciparum* genes using non-redundant oligonucleotides inferred from protein sequences of other organisms, 247
- Jarroll, E.L., Manning, P., Lindmark, D.G., Coggins, J.R. and Erlandsen, S.L.
Giardia cyst wall-specific carbohydrate: evidence for the presence of galactosamine, 121
- Jenkins, M.C., Danforth, H.D., Lillehoj, H.S. and Fetterer, R.H.
cDNA encoding an immunogenic region of a 22 kilodalton surface protein of *Eimeria acervulina* sporozoites, 153
- Karess, R.E., see Button, L.L., 271
- Kelly, C., see Rodrigues, V., 7
- Kelly, C., see Selkirk, M.E., 229

- Kelly, S.L., see Hyde, J.E., 247
 Kemp, D.J., see Favaloro, J.M., 297
 Klein, H.L., see Button, L.L., 271
 Knapp, B., Hundt, E., Nau, U. and Küpper, H.A.
 Molecular cloning, genomic structure and localization in a blood stage antigen of *Plasmodium falciparum* characterized by a serine stretch, 73
 Knight, M., see Rodrigues, V., 7
 Koller, B., see Huber, M., 285
 Krungkrai, J., Webster, H.K. and Yuthavong, Y.
 De novo and salvage biosynthesis of pteroylglutamates in the human malaria parasite, *Plasmodium falciparum*, 25
 Kuhns, J., see Reis, M.G., 113
 Küpper, H.A., see Knapp, B., 73
 Kwan-Lim, G.-E., see Maizels, R.M., 213
- Lensen, T.H.W., see Delves, C.J., 105
 Lillehoj, H.S., see Jenkins, M.C., 153
 Lindmark, D.G., see Jarroll, E.L., 121
- Maizels, R.M., Gregory, W.F., Kwan-Lim, G.-E. and Selkirk, M.E.
 Filarial surface antigens: the major 29 kilodalton glycoprotein and a novel 17-200 kilodalton complex filarial surface antigens from adult *Brugia malayi* parasites, 213
 Maizels, R.M., see Selkirk, M.E., 229
 Mama, K., see Diffley, P., 1
 Manning, P., see Jarroll, E.L., 121
 Marr, J.J., see Goad, L.J., 179
 Marshall, V.M., see Favaloro, J.M., 297
 Matile, H., see Delves, C.J., 105
 McBride, J.S., see Clark, J.T., 15
 McGuire, T.C., see Barbet, A.F., 191
 McMaster, W.R., see Button, L.L., 271
 Meadows, H., see Rodrigues, V., 7
 Medina-Acosta, E., see Button, L.L., 271
 Merritt, Jr., J.W., see Aggarwal, A., 39
 Mirelman, D., see Huber, M., 285
 Myler, P.J., see Barbet, A.F., 191
- Nakamura, Y., see Watanabe, J.-i., 163
 Nash, T.E., see Aggarwal, A., 39
 Nau, U., see Knapp, B., 73
 Nielsen, L., see Selkirk, M.E., 229
 Nonnengässer, C., see Bülow, R., 85
- Ostoa-Saloma, P., Cabrera, N., Becker, I. and Perez-Montfort, R.
 Proteinases of *Entamoeba histolytica* associated with different subcellular fractions, 133
 Overath, P., see Bülow, R., 85
- Paindavoine, P., Zampetti-Bosseler, F., Coquelet, H., Pays, E. and Steinert, M.
 Different allele frequencies in *Trypanosoma brucei brucei* and *Trypanosoma brucei gambiense* populations, 61
 Partono, F., see Selkirk, M.E., 229
 Pays, E., see Paindavoine, P., 61
 Perez-Montfort, R., see Ostoa-Saloma, P., 133
- Perlmann, P., see Franzén, L., 201
 Pettersson, U., see Franzén, L., 201
 Ponnudurai, T., see Delves, C.J., 105
 Prichard, R.K., see Tang, L., 145
- Reis, M.G., Kuhns, J., Blanton, R. and Davis, A.H.
 Localization and pattern of expression of a female specific mRNA in *Schistosoma mansoni*, 113
 Revel, M., see Huber, M., 285
 Ridley, R.G., see Delves, C.J., 105
 Rodrigues, V., Chaudhri, M., Knight, M., Meadows, H., Chambers, A.E., Taylor, W.R., Kelly, C. and Simpson, A.J.G.
 Predicted structure of a major *Schistosoma mansoni* egg-shell protein, 7
 Rozenblatt, S., see Huber, M., 285
 Russell, D.G., see Button, L.L., 271
- Sayers, G., see Selkirk, M.E., 229
 Scaife, J.G., see Delves, C.J., 105
 Schmatz, D.M., Baginsky, W.F. and Turner, M.J.
 Evidence for and characterization of a mannitol cycle in *Eimeria tenella*, 263
 Scholler, J.K., see Aline, Jr., R.F., 169
 Selkirk, M.E., Nielsen, L., Kelly, C., Partono, F., Sayers, G. and Maizels, R.M.
 Identification, synthesis and immunogenicity of cuticular collagens from the filarial nematodes *Brugia malayi* and *Brugia pahangi*, 229
 Selkirk, M.E., see Maizels, R.M., 213
 Simpson, A.J.G., see Rodrigues, V., 7
 Sims, P.F.G., see Hyde, J.E., 247
 Snewin, V.A., see Hyde, J.E., 247
 Snounou, G., see Viriyakosol, S., 93
 Steinert, M., see Paindavoine, P., 61
 Stuart, K., see Aline, Jr., R.F., 169
- Tanabe, K., see Watanabe, J.-i., 163
 Tang, L. and Prichard, R.K.
 Characterization of tubulin from *Brugia malayi* and *Brugia pahangi*, 145
 Taylor, W.R., see Rodrigues, V., 7
 Turner, M.J., see Schmatz, D.M., 263
- Viriyakosol, S., Snounou, G. and Brown, K.N.
 The use of a DNA probe for the differentiation of rodent malaria strains and species, 93
- Wahlgren, M., see Franzén, L., 201
 Wählin, B., see Franzén, L., 201
 Watanabe, J.-i., Hori, H., Tanabe, K. and Nakamura, Y.
 Phylogenetic association of *Pneumocystis carinii* with the 'Rhizopoda/Myxomycota/Zygomycota group' indicated by comparison of 5S ribosomal RNA sequences, 163
 Webster, H.K., see Krungkrai, J., 25
 Wigzell, H., see Franzén, L., 201
 Williams, R.O., see Barbet, A.F., 191
 Wilson, C.F., see Clark, J.T., 15
 Yuthavong, U., see Krungkrai, J., 25
- Zampetti-Bosseler, F., see Paindavoine, P., 61

Subject Index

- Actin, 247
- β -N-Acetylgalactosaminidase, 121
- Amastigote, 179
- Amino hexose, 121
- Antigenic variation, 39, 191
- Asparagine-rich, 201
- Benzimidazole, 145
- Biomphalaria glabrata*, 49
- Brugia*, 229
- Brugia malayi*, 145, 213
- Brugia pahangi*, 145, 213
- Calorimetry, 57
- Carbohydrate, 121
- Carbohydrate metabolism, 263
- Cell cycle, 1
- Chemotherapy, 179
- Chromosome location, 271
- Circular episome, 285
- Codon match index, 247
- Collagen, 229
- Common proteins, 49
- Cross reactivity, 229
- Cuticle, 213
- Cuticle structure, 229
- Cysteine-rich protein, 39
- Cytospectrophotometry, 1
- De novo folate synthesis, 25
- Diagnosis, 105
- Differentiation, 85
- Dihydrofolate reductase-thymidylate synthetase, 247
- DNA hybridisation, 93
- cDNA, 153
- cDNA sequence, 297
- rDNA organization, 285
- 25S rDNA sequence, 285
- Eggshell protein, 7
- Eimeria*, 153
- Eimeria tenella*, 263
- Electron microscopy, 285
- Entamoeba histolytica*, 133, 285
- Epimastigote, 179
- Exon-intron pattern, 73
- Female specific cDNA, 113
- Fluconazole, 179
- Fluorescent probe, 57
- Folate biosynthesis, 25
- Galactosamine, 121
- Gene evolution, 191
- Gene expression, 113
- Gene localization, 113
- Gene organization, 271
- Gene sequence, 7
- Gene structure, 73
- Genetic exchange, 61
- Giardia*, 121
- Giardia lamblia*, 39
- Glycosylation, 213
- Growth rate, 1
- Hexokinase, 263
- Hydrophobic chromatography, 133
- In situ hybridization, 113
- Itraconazole, 179
- Ketoconazole, 179
- Leishmania*, 271
- Major surface glycoprotein, 271
- Malaria, 93, 105, 201
- Mannitol, 263
- Mannitol cycle, 263
- Mannitol dehydrogenase, 263
- Mannitol-1-phosphatase, 263
- Mannitol-1-phosphate dehydrogenase, 263
- Membrane, 57
- Metabolic labeling, 39
- Monoclonal antibody, 15, 201
- Nematode, 145, 213
- Nucleic acid hybridisation, 247
- Octapeptide-repeat antigen, 297
- Oligonucleotide probe, 247
- Peptide map, 213
- Plasmodium berghei*, 93
- Plasmodium chabaudi*, 93
- Plasmodium falciparum*, 15, 25, 73, 105, 201, 297
- Plasmodium vinckei*, 93
- Plasmodium yoelii*, 93
- Pneumocystis carinii*, 163
- Polyols, 263
- Polyribosome, 169
- Predicted structure, 7
- Proteinase, 133
- Protozoan, 39
- Protozoan cyst wall, 121
- Pteroylpentaglutamate, 25
- Qinghaosu, 57
- Recombinant antigen, 153
- Reducing agent, 213
- Repetitive DNA, 105, 201

Restriction fragment length polymorphism, 61
5S Ribosomal RNA, 163
Rodent, 93

Salvage pathway, 25
Schistosoma mansoni, 7, 49
Serine protein, 73
Sporozoite, 153
Sterol, 179
Subcellular fraction, 133
Substrate gel electrophoresis, 133
Surface coat, 85
Surface glycoprotein, 15, 191
Surface iodination, 39
Synthesis, 229

Taxonomy, 93, 163
Transcription, 169, 271
Triton X-114 phase separation, 133
Trypanosoma b. gambiense speciation, 61
Trypanosoma brucei, 85, 169, 191
Trypanosoma brucei gambiense, 1
Trypanosoma cruzi, 179
Trypanosome, 61
Trypomastigote, 179
Tubulin, 247
Tubulin isoform, 145
Turnover, 85

Variant surface glycoprotein, 85, 169
Vitelline cell, 113

